

**CONFIDENTIAL**



**UNIVERSITI TUN HUSSEIN ONN MALAYSIA**

**FINAL EXAMINATION  
SEMESTER I  
SESSION 2013/2014**

COURSE NAME : BASIC PHYSIC 2  
COURSE CODE : DAS 14403  
PROGRAMME : 2 DAE  
EXAMINATION DATE : DECEMBER 2013/JANUARY 2014  
DURATION : 2 ½ HOURS  
INSTRUCTION : ANSWER ALL QUESTIONS IN  
SECTION A AND TWO (2)  
QUESTIONS IN SECTION B

THIS QUESTION PAPER CONSISTS OF NINE (9) PAGES

**CONFIDENTIAL**

## SECTION A

**Q1** (a) A two slits interference experiment is used to determine the unknown wavelength of laser light source. The slits separated by  $2.00 \times 10^{-4}$  m apart and a screen at a distance of 1.00 m. The third bright band out of central bright band is found to be 9.49 mm from the center of the screen. Find

- (i) the definition of interference and give two type of interference.
- (ii) the wavelength of the light.
- (iii) the distance between the  $m = 0$  and  $m = 1$  bright fringes.
- (iv) how far apart would the slit have to be so that the fourth minimum (dark fringe) would occur at 9.49 mm from the center of the screen?

( 12 marks)

(b) A beam of unpolarized light of intensity  $I_0$  passes through a series of ideal polarizing filters with their polarizing directions turned to various angles as shown in **Figure Q1 (b)**. If the light that emerges from the system has an intensity of  $46.0 \text{ Wm}^{-2}$ , use  $I = I_0 \cos^2 \theta$ . Find

- (i) the definition of Malus's law on polarization of light.
- (ii) the intensity of the incident of the light.
- (iii) the light intensity at point C after the middle filter was removed

(13 marks)

- Q2** (a) State the law of reflection and define the index of refraction of a material. (5 marks)
- (b) **Figure Q2 (b)(i)** shows a beam of light incident upon a film of oil at  $\theta_1 = 30^\circ$  from the vertical. The light is refracted in the oil and then refracted again as it enters the water. **Figure Q2 (b)(ii)** shows the light going from water to oil and then totally internal reflecting at the oil / air interface. Determine the
- (i) angle refraction in water (**Figure Q2 (b)(i)**).
- (ii) incident angle in the water (**Figure Q2 (b)(ii)**).
- Given  $n_{\text{water}} = 1.33$ ,  $n_{\text{air}} = 1.00$ ,  $n_{\text{oil}} = 1.48$  (10 marks)
- (c) After leaving some presents under a tree, Santa notices his image in a shiny, spherical Christmas ornament. The ornament is 8.50 cm in diameter and 1.10 m away from Santa as shown in **Figure Q2 (c)**. He curious to know the location and size of his image, Santa consults a book on physics. Knowing that Santa likes to check it twice, what result should he obtain assuming his height is 1.75m? Determine the
- (i) type of mirror.
- (ii) focal length of ornament,  $f$ .
- (iii) image distance,  $d_i$ .
- (iv) magnification,  $m$  of the image.
- (v) image height. (10 marks)

## SECTION B

**Q3 (a)** It is reasonable to assume that the bulk modulus of blood is about the same as that of water. As one goes deeper and deeper in the ocean, the pressure increases by  $1.0 \times 10^4$  Pa for every meter below the surface. The reciprocal of the bulk modulus  $B$  of a liquid is compressibility,  $k$  so  $k = \frac{1}{B}$ . Given the Bulk Modulus of water is 2.2 GPa and 1 atm is equal  $1.013 \times 10^5$  Pa is the pressure exerted by earth's atmosphere at sea level. Find the

- (i) compressibility,  $k$  of bulk modulus of water per atmosphere of pressure (answer in  $\text{atm}^{-1}$ )
- (ii) If a diver goes down 23 m in the ocean, how much each cubic centimeter of her blood change in volume?
- (iii) How deep must a diver go so that each drop of blood compresses to half its volume at the surface?

(12 marks)

(b) When a block of volume  $1.00 \times 10^{-3} \text{ m}^3$  is hung from a spring scale as shown in **Figure Q3 (c)(i)**, the scale reads 10.0 N. When the same block is then placed in an unknown liquid, it floats with  $2/3$  of its volume submerged as suggested in **Figure Q3 (c)(ii)**, The density of water is  $1.00 \times 10^3 \text{ kgm}^{-3}$ . Determine the mass of the block and the density of the unknown liquid

(13 marks)

**Q4 (a)** At  $25^\circ\text{C}$ , a steel sphere bearing is 4.0 cm in diameter is  $6 \times 10^{-4}$  cm larger than the inside of a bronze ring. Determine the single temperature of both ball and ring at which the sphere just slips through the ring.  
[Given the  $\alpha_{\text{Al}} = 11 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$ ,  $\alpha_{\text{Br}} = 19 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$ ]

(13 marks)

(b) A glass flask filled "to the mark" with  $100.0 \text{ cm}^3$  of mercury at  $18^\circ\text{C}$ . If the flask and its contents are heated to  $38^\circ\text{C}$ , find

- (i) how much mercury will be above the mark.
- (ii) if use  $150 \text{ cm}^3$  glass test tube and replace the mercury with water, how much water will be above the mark.

[Given the coefficient of thermal expansion,  $\alpha_{\text{glass}} = 9.0 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$  and  $\gamma_{\text{mercury}} = 182 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$  and  $\gamma_{\text{water}} = 207 \times 10^{-6} \text{ }^\circ\text{C}^{-1}$ ]

(12 marks)


- Q5** (a) A student prepares a standing waves experiment. He is using a metal string under a tension of 88.2 N. Its length is 50.0 cm and its mass is  $5 \times 10^{-4}$  kg. When the vibrator is turned on the string is found to rapidly develop a large, stable transverse standing waves consisting of four equal sections. Determine
- (i) the velocity for transverse waves on the string.
  - (ii) the frequencies of its fundamental, first overtone, second overtone.
- (10 marks)
- (b) A source transmitted sound wave with an output power 3 Watt. Calculate the
- (i) sound intensity at a distance 6.5 m from the source.
  - (ii) intensity level of the sound heard by the observer.
  - (iii) for the maximum sound intensity level in the workplace is 90.0 dB. Within one factory, 32 identical machines produce a sound intensity level of 92.0 dB. How many machines must be removed to ensure the sound intensity must not exceed the maximum?
- (15 marks)

**- END OF QUESTION -**


**FINAL EXAMINATION**

SEMESTER / SESSION : SEM I / 2013/2014  
 COURSE : BASIC PHYSICS 2


PROGRAMME : 3 DAC  
 COURSE CODE : DAS 14403




**FIGURE Q1 (b)**



**FIGURE Q2(b)(i)**



**FIGURE Q2(b)(ii)**




**FIGURE Q2 (c)**

**FINAL EXAMINATION**

SEMESTER / SESSION : SEM I / 2013/2014  
COURSE : BASIC PHYSICS 2

PROGRAMME : 3 DAC  
COURSE CODE : DAS 14403



**FIGURE Q3(b)(i) in water and (ii) in unknown liquid.**

**FINAL EXAMINATION**SEMESTER / SESSION : SEM I / 2013/2014  
COURSE : BASIC PHYSICS 2PROGRAMME : 3 DAC  
COURSE CODE : DAS 14403**LIST OF CONSTANTS**

1. Gravity acceleration,  $g = 9.81 \text{ m/s}^2$
2. Speed of light in air,  $c = 3 \times 10^8 \text{ m/s}$
3. Speed of sound,  $v_{\text{sound}} = 335 \text{ m/s}$
4. Threshold of sound intensity,  $I_o = 1 \times 10^{-12} \text{ W/m}^2$
5. Atmospheric pressure,  $P_{\text{atm}} = 1.0 \times 10^5 \text{ Pa}$
6. Specific heat of water,  $c_{\text{water}} = 4.186 \text{ kJkg}^{-1}\text{K}^{-1}$
7. Specific heat of copper,  $c_{\text{copper}} = 0.385 \text{ kJkg}^{-1}\text{K}^{-1}$
8. Specific heat of lead,  $c_{\text{lead}} = 0.13 \text{ kJkg}^{-1}\text{K}^{-1}$
8. Latent heat of fusion of water,  $L_f = 333.7 \times 10^3 \text{ J/kg}$
10. Latent heat of vaporization of water,  $L_v = 2256 \times 10^3 \text{ J/kg}$
11. Density of seawater,  $\rho_{\text{seawater}} = 1030 \text{ kg/m}^3$
12. Density of water,  $\rho_{\text{water}} = 1000 \text{ kg/m}^3$
13. Refractive index ( $n_{\text{air}}$ ) of air = 1.00
14. Refractive index ( $n_{\text{water}}$ ) of water = 1.333
15. Refractive index ( $n_{\text{ice}}$ ) of ice = 1.304



**FORMULAE**

SEMESTER / SESSION : SEM I / 2013/2014  
 COURSE : BASIC PHYSICS 2

PROGRAMME : 3 DAC  
 COURSE CODE : DAS 14403

$y = \frac{(m + \frac{1}{2})\lambda D}{d}$	$\sin\theta = \frac{m\lambda}{d}$	$y = \frac{m\lambda L}{d}$
$T_F = 1.8T_C + 32^\circ F$	$\Delta L = \alpha L_o \Delta T$	$\Delta A = \beta A_o \Delta T$
$Q = mc\Delta T$	$Q = mL_f$	$Q = mL_v$
$\Delta V = \gamma V_o \Delta \theta$	$\gamma = \gamma_{\text{apparent}} + \gamma_{\text{glass}}$	$I = I_o \cos^2 \theta$
$A_{\text{circle}} = \pi r^2$	$F_{\text{net}} = W - F_B$	$A_{\text{sphere}} = 4\pi r^2$
$P = \rho gh$	$F = \rho gV$	$n_1 \sin\theta_1 = n_2 \sin\theta_2$
$\mu = \rho \pi r^2$	$V_{\text{sphere}} = \frac{4}{3} \pi r^3$	$\frac{Q}{t} = \frac{\Delta T}{\sum R_n}$
$\lambda = \frac{v}{f}$	$\mu = \frac{m}{l}$	$\beta = 10 \log \left( \frac{I}{I_o} \right)$
$m = -\frac{d_i}{d_o}$	$f_o = f_s \frac{(v \pm v_o)}{(v \mp v_s)}$	$\frac{1}{d_o} + \frac{1}{d_i} = \frac{1}{f}$
$\frac{Q}{t} = \kappa A \frac{\Delta T}{d}$	$\frac{Q}{t} = \kappa A \left( \frac{T_{\text{hot}} - T_{\text{cold}}}{d} \right)$	$T = \left( \frac{X_T - X_o}{X_{100} - X_o} \right) \times 100^\circ C$
$P = \frac{F}{A}$	$\rho = \frac{m}{V}$	$\Delta P = \frac{F_1}{A_1} = \frac{F_2}{A_2}$