

**CONFIDENTIAL**



**UNIVERSITI TUN HUSSEIN ONN MALAYSIA**

**FINAL EXAMINATION  
SEMESTER I  
SESSION 2019/2020**

COURSE NAME : ENGINEERING MATHEMATICS II  
COURSE CODE : DAM 21303 / DAE 23403  
PROGRAMME CODE : DAM / DAE  
EXAMINATION DATE : DECEMBER 2019 / JANUARY 2020  
DURATION : 3 HOURS  
INSTRUCTION : ANSWERS ALL QUESTIONS IN SECTION A AND ANSWER **THREE (3)** QUESTIONS IN SECTION B

THIS QUESTION PAPER CONSISTS OF **EIGHT (8)** PAGES

**TERBUKA**

**CONFIDENTIAL**

**SECTION A**

**Q1** Solve the following equations by using the method of undetermined coefficient.

(a)  $\frac{d^2y}{dx^2} - 3\frac{dy}{dx} - 4y = 2 \cos 2x.$

(8 marks)

(b)  $\frac{d^2y}{dx^2} - 4y = x + 5 \sin 3x$  ;  $y(0) = 2$  and  $y'(0) = 0.$

(12 marks)

**Q2** By using the method of variation of parameters to find the general solution of the given second order differential equation.

(a)  $y'' - 6y' + 9y = (3 - x)e^{3x}.$

(10 marks)

(b)  $\frac{d^2y}{dx^2} - 7\frac{dy}{dx} + 10y = -\frac{x}{\pi} - 3$

(10 marks)

**SECTION B**

**Q3** (a) Find the integrals of

(i)  $\int (8x^3 + 3e^{-5x}) dx.$

(4 marks)

(ii)  $\int \left( \cos 7x - \frac{4}{(3x+1)} \right) dx.$

(4 marks)

(iii)  $\int x^2 \ln x dx.$

(5 marks)

(b) Find the approximate value for  $\int_0^1 \sqrt{x^2 + 1} dx$  using trapezoidal rule by taking step size,  $h = 0.2.$

(7 marks)

**TERBUKA**

**Q4** (a) Given two curves  $y = \sqrt{x-1}$  and  $y = (x-1)^2$ .

(i) Sketch the graphs of the curves. (5 marks)

(ii) By using the cylindrical shells method, find the volume of the solid generated when the bounded region is rotated about  $x$ -axis. (8 marks)

(b) Find the arc length of the curve  $y = \frac{4}{3}x^{\frac{3}{2}} - 1$  when  $1 \leq x \leq 2$ . (7 marks)

**Q5** (a) Find the solution of the given linear differential equation.

$$(x+2)\frac{dy}{dx} + y = (x+2)e^{2x}, \quad \text{IVP: } y(0) = 2.$$

(10 marks)

(b) Given the first order differential equation

$$(y - xy^2 + 2ye^x)dx + (x - x^2y + 2e^x)dy = 0.$$

(i) Show that the equation is an exact ordinary differential equation. (2 marks)

(ii) Find the general solution of the equation. (8 marks)

**Q6** (a) As a worker in a factory, you need to remove a heavy metal with its core temperature of  $1000^\circ\text{F}$  from a furnace and placed the metal on a table in a room that had a constant temperature of  $72^\circ\text{F}$ . One hour after it is removed the core temperature is  $910^\circ\text{F}$ . The temperature of the metal must be below  $540^\circ\text{F}$  before you can transfer it to the next section.

(i) Given  $\frac{dT}{T - T_s} = -kdt$ . Show that  $T - T_s = Ae^{-kt}$ . (4 marks)

(ii) By using  $T - T_s = Ae^{-kt}$ , with  $T(0) = 1000^\circ\text{F}$  and  $T_s = 72^\circ\text{F}$ , find the constant  $A$ . Hence find  $T(t)$ . (4 marks)

- (iii) Given the observed temperatures of the metal, given  $T(1) = 910^\circ\text{F}$ , find the constant  $k$ . (4 marks)
- (iv) Find the time taken for the temperature of the metal to be below  $540^\circ\text{F}$ . (3 marks)
- (b) The world population growth is described by  $y(t) = y_0 e^{k(t-t_0)}$  with  $t$  measured in years.
  - (i) If the population increased 2011 by 3% from 2010 to 2011, find  $k$ . (3 marks)
  - (ii) If the population in  $t_0 = 2010$  was 5 million people, find the actual population for 2020 predicted by the given equation. (2 marks)

**Q7** (a) By using the integration by parts to show that  $\int_0^2 \frac{1}{\sqrt{x}} e^{\frac{1}{2}x} dx = 2\sqrt{x}e^{\frac{1}{2}x} - \int_0^2 \sqrt{x}e^{\frac{1}{2}x} dx$ . (3 marks)

(b) By using Simpson's Rule with  $h = 0.4$ , find the approximate value of  $\int_1^3 \frac{3x}{x+5} dx$ . (5 marks)

- (c) The equations of two curves are given by  $y = x^2 - 1$  and  $y = \frac{6}{x^2}$ .
- (i) Sketch the two curves on the same coordinates axes. (3 marks)
  - (ii) Find the coordinates of the points of intersection of the two curves. (3 marks)
  - (iii) Calculate the volume of the solid formed when the region bounded by the two curves and the line  $x = 1$  is revolved completely about the  $y$ -axis. (6 marks)

- END OF QUESTIONS -

TERBUKA

**FINAL EXAMINATION**

SEMESTER / SESSION : SEM I / 2019/2020  
 COURSE NAME : ENGINEERING MATHEMATICS II

PROGRAMME CODE : DAM / DAE  
 COURSE CODE : DAM 21303 / DAE23403

**Formula**

**Table 1: Characteristic Equation and General Solution**

Homogeneous Differential Equation: $ay'' + by' + cy = 0$		
Characteristics Equation: $am^2 + bm + c = 0$		
$m = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$		
Case	Roots of Characteristics Equation	General Solution
1	Real and Distinct: $m_1 \neq m_2$	$y_h(x) = Ae^{m_1x} + Be^{m_2x}$
2	Real and Equal: $m_1 = m_2 = m$	$y_h(x) = (A + Bx)e^{mx}$
3	Complex Roots: $m = \alpha \pm i\beta$	$y_h(x) = e^{\alpha x}(A \cos \beta x + B \sin \beta x)$

**Table 2: Particular Solution of Nonhomogeneous Equation**

$ay'' + by' + cy = f(x)$

$f(x)$	$y_p(x)$
$P_n(x) = A_n x^n + A_{n-1} x^{n-1} + \dots + A_1 x + A_0$	$x^r (B_n x^n + B_{n-1} x^{n-1} + \dots + B_1 x + B_0)$
$Ce^{\alpha x}$	$x^r (Pe^{\alpha x})$
$C \cos \beta x$ or $C \sin \beta x$	$x^r (P \cos \beta x + Q \sin \beta x)$
$P_n(x)e^{\alpha x}$	$x^r (B_n x^n + B_{n-1} x^{n-1} + \dots + B_1 x + B_0)e^{\alpha x}$
$P_n(x) \begin{cases} \cos \beta x \\ \sin \beta x \end{cases}$ or	$x^r (B_n x^n + B_{n-1} x^{n-1} + \dots + B_1 x + B_0) \cos \beta x$ $+ x^r (B_n x^n + B_{n-1} x^{n-1} + \dots + B_1 x + B_0) \sin \beta x$

**Notes:**  $r$  is the smallest non negative integer to ensure no alike term between  $y_p(x)$  and  $y_h(x)$ .

**TERBUKA**



**FINAL EXAMINATION**

SEMESTER / SESSION : SEM I / 2019/2020  
 COURSE NAME : ENGINEERING MATHEMATICS II

PROGRAMME CODE : DAM / DAE  
 COURSE CODE : DAM 21303 / DAE23403

**Table3: Variation of Parameters Method**

Homogeneous solution, $y_h(x) = Ay_1 + By_2$	
Wronskian function, $W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix} = y_1 y_2' - y_2 y_1'$	
$u_1 = -\int \frac{y_2 f(x)}{aW} dx + A$	$u_2 = \int \frac{y_1 f(x)}{aW} dx + B$
General solution, $y(x) = u_1 y_1 + u_2 y_2$	

**Table 4: Trigonometry Identities**

$\sin^2 t + \cos^2 t = 1$
$\sin^2 t = \frac{1}{2}(1 - \cos 2t)$
$\cos^2 t = \frac{1}{2}(1 + \cos 2t)$

**Table 5: Partial Fraction**

$\frac{a}{(s+b)(s-c)} = \frac{A}{s+b} + \frac{B}{s-c}$
$\frac{a}{s(s-b)(s-c)} = \frac{A}{s} + \frac{B}{s-b} + \frac{C}{s-c}$
$\frac{a}{(s+b)^2} = \frac{A}{s+b} + \frac{B}{(s+b)^2}$
$\frac{a}{(s+b)(s^2+c)} = \frac{A}{s+b} + \frac{Bs+C}{s^2+c}$

**TERBUKA**

**FINAL EXAMINATION**

SEMESTER / SESSION : SEM I / 2019/2020  
 COURSE NAME : ENGINEERING MATHEMATICS II

PROGRAMME CODE : DAM / DAE  
 COURSE CODE : DAM 21303 / DAE23403

**Table 6: Integration and Differentiation**

Integration	Differentiation
$\int x^n dx = \frac{x^{n+1}}{n+1} + C$	$\frac{d}{dx} x^n = nx^{n-1}$
$\int \frac{1}{x} dx = \ln x  + C$	$\frac{d}{dx} \ln x = \frac{1}{x}$
$\int \frac{1}{a-bx} dx = -\frac{1}{b} \ln a-bx  + C$	$\frac{d}{dx} \ln(ax+b) = \frac{a}{ax+b}$
$\int e^{ax} dx = \frac{1}{a} e^{ax} + C$	$\frac{d}{dx} e^{ax} = ae^{ax}$
$\int \sin ax dx = -\frac{1}{a} \cos ax + C$	$\frac{d}{dx} \sin ax = a \cos ax$
$\int \cos ax dx = \frac{1}{a} \sin ax + C$	$\frac{d}{dx} \cos ax = -a \sin ax$
$\int \sec^2 x dx = \tan x + C$	$\frac{d}{dx} \tan x = \sec^2 x$
$\int \csc^2 x dx = -\cot x + C$	$\frac{d}{dx} \cot x = -\csc^2 x$
$\int u dv = uv - \int v du$	$\frac{d}{ds} (uv) = u \frac{dv}{ds} + v \frac{du}{ds}$
$\int_a^b f(x) dx = F(b) - F(a)$	$\frac{d}{ds} \left( \frac{u}{v} \right) = \frac{v \frac{du}{ds} - u \frac{dv}{ds}}{v^2}$

TERBUKA

**FINAL EXAMINATION**

SEMESTER / SESSION : SEM I / 2019/2020  
 COURSE NAME : ENGINEERING MATHEMATICS II

PROGRAMME CODE : DAM / DAE  
 COURSE CODE : DAM 21303 / DAE23403

**Area of Region**

$$A = \int_a^b [f(x) - g(x)] dx \quad \text{or} \quad A = \int_c^d [w(y) - v(y)] dy$$

**Volume Cylindrical Shells**

$$V = \int_a^b 2\pi x f(x) dx \quad \text{or} \quad V = \int_c^d 2\pi y f(y) dy$$

**Arc Length**

$$L = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx \quad \text{or} \quad L = \int_c^d \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

**Simpson's Rule**

$$\int_a^b f(x) dx \approx \frac{h}{3} \left[ (f(a) + f(b)) + 4 \sum_{\substack{i=1 \\ i \text{ odd}}}^{n-1} f(a+ih) + 2 \sum_{\substack{i=2 \\ i \text{ even}}}^{n-2} f(a+ih) \right]; \quad n = \frac{b-a}{h}$$

**Trapezoidal Rule**

$$\int_a^b f(x) dx \approx \frac{h}{2} \left[ f(a) + f(b) + 2 \sum_{i=1}^{n-1} f(a+ih) \right]; \quad n = \frac{b-a}{h}$$

**TERBUKA**