



UNIVERSITI TUN HUSSEIN ONN MALAYSIA

FINAL EXAMINATION SEMESTER I SESSION 2015/2016

COURSE NAME : ENGINEERING MATHEMATICS V
COURSE CODE : BEE 31702 / BWM 20502
PROGRAMME : BACHELOR OF ELECTRONIC
ENGINEERING WITH HONOURS/
BACHELOR OF ELECTRICAL
ENGINEERING WITH HONOURS
EXAMINATION DATE : DECEMBER 2015 / JANUARY 2016
DURATION : 2 HOURS 30 MINUTES
INSTRUCTION : ANSWER ALL QUESTIONS

THIS QUESTION PAPER CONSISTS OF TEN (10) PAGES

CONFIDENTIAL

- Q1** In a study of pollution in a water stream, the concentration of pollution is measured at 5 different locations. The locations are at different distances to the pollution source. These distances and the average pollution are given in **Table 1**.

TABLE 1

Distance to the pollution (in km)	Average concentration (avg)
10	9.3
8	9.675
6	10.25
4	10.225
2	11.5

- (a) Produce the equation of the least squares line that will enable us to predict the average concentration in terms of the distance to the pollution. Interpret the results. (12 marks)
- (b) Identify the average concentration when the relative distance is 12 km. (2 marks)
- (c) Find and interpret the Pearson correlation coefficient. (6 marks)

- Q2** A recent study is performed to discover whether gender affects education level. The study surveyed a random sample of 395 people and each person was asked to report the highest education level they obtained. The data that resulted from the survey is summarized in **Table 2**. The claim from the study suggested that the means for both genders who achieved greater education would no longer be 50.

TABLE 2

	High School	Bachelors	Masters	Ph. D	Total
Female	60	54	46	41	201
Male	40	44	53	57	194
Total	100	98	99	98	395

- (a) Find the means and sample deviations for both female and male samples. (4 marks)
- (b) State the null and alternative hypothesis. (2 marks)
- (c) Identify the Type I and Type II errors that corresponds to the hypothesis above. (2 marks)

CONFIDENTIAL

- (d) Test this claim using the critical value approach with 5 % significance for both males and females separately. (12 marks)

Q3 A general hospital recorded that 75% of the cancer patients are died after the chemotherapy treatment. Assume that the distribution of the cancer patients who died after the chemotherapy treatment is binomially distributed. If 5 patients are selected randomly,

- (a) Find the mean, variance and standard deviation of the distribution. (4 marks)
- (b) Calculate the probability that only two of the patients are recovered. (2 marks)
- (c) Produce an approximate Poisson distribution probability to find at most two patients are died. (5 marks)
- (d) Produce an approximate normal probability distribution to find exactly 36 patients died after the chemotherapy, if 6000 patients are randomly selected and the probability of success of the cancer patients are died is 0.005. (9 marks)

Q4 (a) According to the following blood types and possible alleles:

- Type O: Must be OO
- Type A: Could be AA or OA
- Type B: Could be BB or OB
- Type AB: Must be AB

Suppose father has OO (type O) and mother has OB (type B). They have 3 children. Each child equally likely to inherit blood type O or type B. Let x is the number of blood type B.

- (i) Identify the sample space of x for three births. (2 marks)
- (ii) Find the probability of x by using p.d.f table. (2 marks)
- (iii) Produce the cumulative distribution function for x . (5 marks)

CONFIDENTIAL

- (b) The lead concentration in gasoline currently ranges from 0.1 to 0.5 grams per litre with probability density function given by

$$f(x)=\begin{cases} 12.5x-1.25 & , \quad 0.1 \leq x \leq 0.5 \\ 0 & , \quad otherwise \end{cases}$$

(i) Find the expected value of x .
(5 marks)

(ii) Calculate the variance of x .
(6 marks)

- Q5** In order to understand the reliability of the internet service, a team from a local Internet Service Provider (ISP), Unifi™, has conducted a study on data transmissions over rural and city areas. The data collected are both taken for a period of one week and is tabulated in **Table 3** below.

TABLE 3

	Rural Area	City Area
Sample size	11	13
Sample mean	2.16	4.82
Sample standard deviation	1.17	1.01

- (a) Identify the mean and standard deviation of the sample mean if the mean of the city area is greater than the mean of the rural area.
(3 marks)
- (b) Find the Z-distribution probability of the city area if the sample mean is between 2.35 and 2.55, exclusive.
(7 marks)
- (c) Calculate the true variance for a 95% confidence interval of the city area using the χ^2 -distribution.
(5 marks)
- (d) Calculate the ratio for both true variances of the areas with a given 90% confidence interval.
(5 marks)

- END OF QUESTIONS -

FINAL EXAMINATION

SEMESTER / SESSION : SEM I / 2014/2015
COURSE : ENGINEERING
MATHEMATICS V

PROGRAMME : 2 BEV / BEJ
COURSE CODE : BEE 31702

Formula

Random Variable :

$$\sum_{i=-\infty}^{\infty} P(x_i) = 1, \quad E(X) = \sum_{\forall x} x \cdot P(x), \quad E(X^2) = \sum_{\forall x} x^2 \cdot P(x), \quad \int_{-\infty}^{\infty} f(x) dx = 1,$$

$$E(X) = \int_{-\infty}^{\infty} x \cdot P(x) dx, \quad E(X^2) = \int_{-\infty}^{\infty} x^2 \cdot P(x) dx, \quad Var(X) = E(X^2) - [E(X)]^2.$$

$$X \sim P_0(\mu), Z = \frac{X - \mu}{\sigma},$$

Sampling Distributions :

$$\overline{xy} = y(-2/1) \equiv \overline{X}$$

Estimations : $X \sim N(\mu, \sigma^2/n)$, $Z = \frac{X - \mu}{\sigma/\sqrt{n}} \sim N(0, 1)$, $T = \frac{\bar{X}_1 - \bar{X}_2}{s/\sqrt{n}}$, $\bar{X}_1 - \bar{X}_2 \sim N\left(\mu_1 - \mu_2, \frac{1}{n_1} + \frac{1}{n_2}\right)$.

(7) —

$$n = \left(\frac{Z_{\alpha/2} \cdot o}{E} \right)^2, \quad \left(\bar{x}_1 - \bar{x}_2 \right) - Z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}} < \mu_1 - \mu_2 < \left(\bar{x}_1 - \bar{x}_2 \right) + Z_{\alpha/2} \sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}},$$

$$\left(\bar{x}_1 - \bar{x}_2 \right) - Z_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} < \mu_1 - \mu_2 < \left(\bar{x}_1 - \bar{x}_2 \right) + Z_{\alpha/2} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}},$$

$$\left(\bar{x}_1 - \bar{x}_2 \right) - t_{\alpha/2,v} \cdot S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}} < \mu_1 - \mu_2 < \left(\bar{x}_1 - \bar{x}_2 \right) + t_{\alpha/2,v} \cdot S_p \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}$$

where Pooled estimate of variance, $S_p^2 = \frac{(n_1-1)s_1^2 + (n_2-1)s_2^2}{n_1+n_2-2}$ with $v = n_1 + n_2 - 2$,

$$\left(\bar{x}_1 - \bar{x}_2 \right) - t_{\alpha/2, v} \sqrt{\frac{1}{n} \left(s_1^2 + s_2^2 \right)} < \mu_1 - \mu_2 < \left(\bar{x}_1 - \bar{x}_2 \right) + t_{\alpha/2, v} \sqrt{\frac{1}{n} \left(s_1^2 + s_2^2 \right)} \text{ with } v = 2(n-1)$$

$$\left(\bar{x}_1 - \bar{x}_2 \right) - t_{\alpha/2, v} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} < \mu_1 - \mu_2 < \left(\bar{x}_1 - \bar{x}_2 \right) + t_{\alpha/2, v} \sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}} \quad \text{with} \quad v = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2} \right)^2}{\frac{\left(s_1^2 \right)^2}{n_1} + \frac{\left(s_2^2 \right)^2}{n_2}},$$

$$\frac{(n-1) \cdot s^2}{\chi_{\alpha/2,v}^2} < \sigma^2 < \frac{(n-1) \cdot s^2}{\chi_{1-\alpha/2,v}^2} \text{ with } v = n-1,$$

$$\frac{s_1^2}{s_2^2} \cdot \frac{1}{f_{\alpha/2}, v_1, v_2} < \frac{\sigma_1^2}{\sigma_2^2} < \frac{s_1^2}{s_2^2} \cdot f_{\alpha/2}, v_2, v_1 \text{ with } v_1 = n_1 - 1 \text{ and } v_2 = n_2 - 1.$$

FINAL EXAMINATION

SEMESTER / SESSION : SEM I / 2014/2015
 COURSE : ENGINEERING
 MATHEMATICS V

PROGRAMME : 2 BEV / BEJ
 COURSE CODE : BEE 31702

Hypothesis Testings :

$$Z = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}}, T = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{S_p \cdot \sqrt{\frac{1}{n_1} + \frac{1}{n_2}}} \text{ with } v = n_1 + n_2 - 2,$$

$$Z = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}}, T = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{1}{n}(s_1^2 + s_2^2)}}, T = \frac{(\bar{X}_1 - \bar{X}_2) - (\mu_1 - \mu_2)}{\sqrt{\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}}} \text{ with}$$

$$v = \frac{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)^2}{\left(\frac{s_1^2}{n_1}\right)^2 + \left(\frac{s_2^2}{n_2}\right)^2}, S_p^2 = \frac{(n_1-1)s_1^2 + (n_2-1)s_2^2}{n_1+n_2-2}, \chi^2 = \frac{(n-1)s^2}{\sigma^2}$$

Simple Linear Regressions :

$$S_{xy} = \sum x_i y_i - \frac{\sum x_i \cdot \sum y_i}{n}, S_{xx} = \sum x_i^2 - \frac{(\sum x_i)^2}{n}, S_{yy} = \sum y_i^2 - \frac{(\sum y_i)^2}{n}, \bar{x} = \frac{\sum x}{n}, \bar{y} = \frac{\sum y}{n},$$

$$\hat{\beta}_1 = \frac{S_{xy}}{S_{xx}}, \hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}, \hat{y} = \hat{\beta}_0 + \hat{\beta}_1 x, r = \frac{S_{xy}}{\sqrt{S_{xx} \cdot S_{yy}}}, SSE = S_{yy} - \hat{\beta}_1 S_{xy}, MSE = \frac{SSE}{n-2},$$

$$T = \frac{\hat{\beta}_1 - \beta_1^*}{\sqrt{\frac{MSE}{S_{xx}}}} \sim t_{n-2}, T = \frac{\hat{\beta}_0 - \beta_0^*}{\sqrt{MSE \left(\frac{1}{n} + \frac{\bar{x}^2}{S_{xx}} \right)}} \sim t_{n-2}.$$

CONFIDENTIAL**FINAL EXAMINATION**


SEMESTER / SESSION : SEM I / 2014/2015
 COURSE : ENGINEERING
 MATHEMATICS V

PROGRAMME : 2 BEV / BEJ
 COURSE CODE : BEE 31702

Basic Distribution and Significance Table**Table 3****AREAS IN TAIL OF THE NORMAL DISTRIBUTION**

The function tabulated is $1 - \Phi(u)$ where $\Phi(u)$ is the cumulative distribution function of a standardised Normal variable u . Thus $1 - \Phi(u) = \frac{1}{\sqrt{2\pi}} \int_u^\infty e^{-u^2/2} du$ is the probability that a

standardised Normal variable selected at random will be greater than a value of u ($= \frac{x-\mu}{\sigma}$)



$\frac{(x - \mu)}{\sigma}$.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641
0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
1.5	.0668	.0655	.0643	.0630	.0618	.0606	.0594	.0582	.0571	.0559
1.6	.0548	.0537	.0526	.0516	.0505	.0495	.0485	.0475	.0465	.0455
1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
2.0	.02275	.02222	.02169	.02118	.02068	.02018	.01970	.01923	.01876	.01831
2.1	.01786	.01743	.01700	.01659	.01618	.01578	.01539	.01500	.01463	.01426
2.2	.01390	.01355	.01321	.01287	.01255	.01222	.01191	.01160	.01130	.01101
2.3	.01072	.01044	.01017	.00990	.00964	.00939	.00914	.00889	.00866	.00842
2.4	.00820	.00798	.00776	.00755	.00734	.00714	.00695	.00676	.00657	.00639
2.5	.00621	.00604	.00587	.00570	.00554	.00539	.00523	.00508	.00494	.00480
2.6	.00466	.00453	.00440	.00427	.00415	.00402	.00391	.00379	.00368	.00357
2.7	.00347	.00336	.00326	.00317	.00307	.00298	.00289	.00280	.00272	.00264
2.8	.00256	.00248	.00240	.00233	.00226	.00219	.00212	.00205	.00199	.00193
2.9	.00187	.00181	.00175	.00169	.00164	.00159	.00154	.00149	.00144	.00139
3.0	.00135									
3.1	.00097									
3.2	.00069									
3.3	.00048									
3.4	.00034									
3.5	.00023									
3.6	.00016									
3.7	.00011									
3.8	.00007									
3.9	.00005									
4.0	.00003									

FINAL EXAMINATION

SEMESTER / SESSION : SEM I / 2014/2015
 COURSE : ENGINEERING
 MATHEMATICS V

PROGRAMME : 2 BEV / BEJ
 COURSE CODE : BEE 31702

Table 7


PERCENTAGE POINTS OF THE t DISTRIBUTION

The table gives the value of $t_{\alpha; \nu}$ — the 100 α percentage point of the t distribution for ν degrees of freedom.

The values of t are obtained by solution of the equation:-

$$\alpha = \Gamma\left(\frac{1}{2}(\nu+1)\right) \left\{ \Gamma\left(\frac{1}{2}\nu\right) \right\}^{-1} (\nu\pi)^{-1/2} \int_0^{\infty} (1+x^2/\nu)^{-(\nu+1)/2} dx$$

Note. The tabulation is for one tail only i.e. for positive values of t . For $|t|$ the column headings for α must be doubled.



$\alpha =$	0.10	0.05	0.025	0.01	0.005	0.001	0.0005
$\nu = 1$	3.078	6.314	12.706	31.821	63.657	318.31	636.62
2	1.886	2.920	4.303	6.965	9.925	22.326	31.598
3	1.638	2.353	3.182	4.541	5.841	10.213	12.924
4	1.533	2.132	2.776	3.747	4.604	7.173	8.610
5	1.476	2.015	2.571	3.365	4.032	5.893	6.869
6	1.440	1.943	2.447	3.143	3.707	5.208	5.959
7	1.415	1.895	2.365	2.998	3.499	4.785	5.408
8	1.397	1.860	2.306	2.896	3.355	4.501	5.041
9	1.383	1.833	2.262	2.821	3.250	4.297	4.781
10	1.372	1.812	2.228	2.764	3.169	4.144	4.587
11	1.363	1.796	2.201	2.718	3.106	4.025	4.437
12	1.356	1.782	2.179	2.681	3.055	3.930	4.318
13	1.350	1.771	2.160	2.650	3.012	3.852	4.221
14	1.345	1.761	2.145	2.624	2.977	3.787	4.140
15	1.341	1.753	2.131	2.602	2.947	3.733	4.073
16	1.337	1.746	2.120	2.583	2.921	3.686	4.015
17	1.333	1.740	2.110	2.567	2.898	3.646	3.965
18	1.330	1.734	2.101	2.552	2.858	3.610	3.922
19	1.328	1.729	2.093	2.539	2.861	3.579	3.883
20	1.325	1.725	2.086	2.528	2.845	3.552	3.850
21	1.323	1.721	2.080	2.518	2.831	3.527	3.819
22	1.321	1.717	2.074	2.508	2.819	3.505	3.792
23	1.319	1.714	2.069	2.500	2.807	3.485	3.767
24	1.318	1.711	2.064	2.492	2.797	3.467	3.745
25	1.316	1.708	2.060	2.485	2.787	3.450	3.725
26	1.315	1.706	2.056	2.479	2.779	3.435	3.707
27	1.314	1.703	2.052	2.473	2.771	3.421	3.690
28	1.313	1.701	2.048	2.467	2.763	3.408	3.674
29	1.311	1.699	2.045	2.462	2.756	3.396	3.659
30	1.310	1.697	2.042	2.457	2.750	3.385	3.646
40	1.303	1.684	2.021	2.423	2.704	3.307	3.551
60	1.296	1.671	2.000	2.390	2.660	3.232	3.460
120	1.289	1.658	1.980	2.358	2.617	3.160	3.373
∞	1.282	1.645	1.960	2.326	2.576	3.090	3.291

FINAL EXAMINATION

SEMESTER / SESSION : SEM I / 2014/2015
 COURSE : ENGINEERING
 MATHEMATICS V
 PROGRAMME : 2 BEV / BEJ
 COURSE CODE : BEE 31702

Table 8

Table of $\chi^2_{\alpha, v}$ – the 100 α percentage point of the χ^2 distribution for v degrees of freedom



$\alpha =$.995	.99	.98	.975	.95	.90	.80	.75	.70	.50	.30	.25	.20	.10	.05	.025	.02	.01	.005	.001	$\alpha = \alpha$
$v = 1$	0.4363	0.0157	0.0628	0.0982	0.0993	0.058	0.0442	0.102	0.148	0.455	1.074	1.323	1.642	2.706	3.841	5.024	5.412	6.633	7.879	10.827	$v = 1$
2	0.060	0.0201	0.0404	0.0506	0.0503	0.211	0.198	0.375	0.713	1.355	2.408	2.773	3.219	4.605	5.991	7.378	7.824	9.210	10.507	13.815	2
3	0.0717	0.115	0.185	0.218	0.252	0.584	0.605	1.213	1.424	2.366	3.665	4.108	4.642	5.251	7.811	9.340	9.837	11.343	12.838	15.288	3
4	0.207	0.429	0.484	0.711	1.064	1.049	1.923	2.195	3.557	4.878	5.385	5.989	7.179	9.488	11.143	11.668	13.277	14.850	18.465	4	
5	0.412	0.551	0.752	0.831	1.145	1.610	2.443	2.675	3.000	4.351	6.064	6.626	7.289	9.236	11.070	12.832	13.388	15.086	16.750	20.517	5
6	0.676	0.872	1.134	1.237	1.635	2.204	3.070	3.455	3.828	5.348	7.231	7.841	8.558	10.645	12.592	14.449	15.033	16.612	18.548	22.457	6
7	0.989	1.239	1.584	1.990	2.187	2.813	3.822	4.255	4.671	6.346	8.383	9.037	9.808	12.017	14.057	16.013	16.622	18.475	20.278	24.322	7
8	1.344	1.646	2.032	2.180	2.733	3.490	4.594	5.071	5.524	7.344	9.524	10.219	11.030	13.362	15.507	17.335	18.168	20.090	21.955	26.125	8
9	1.705	2.083	2.532	2.700	3.325	4.168	5.389	5.899	6.393	8.343	10.656	11.389	12.242	14.684	16.919	19.023	19.679	21.666	22.589	27.377	9
10	2.156	2.558	3.059	3.247	3.940	4.855	6.179	6.737	7.267	9.342	11.781	12.549	13.412	15.987	18.307	20.463	21.161	23.209	25.188	29.538	10
11	2.603	3.050	3.609	3.818	4.575	5.378	6.989	7.584	8.148	10.341	12.899	13.701	14.631	17.275	19.575	21.920	22.618	24.725	26.757	31.384	11
12	3.074	3.571	4.178	4.404	5.226	6.304	7.907	8.438	9.034	11.340	14.014	14.845	15.812	18.549	21.026	23.337	24.054	26.217	28.300	32.909	12
13	3.565	4.107	4.765	5.009	5.892	7.042	8.634	9.299	9.926	12.340	15.119	15.984	16.955	19.812	22.362	24.736	25.472	27.688	29.819	34.528	13
14	4.075	4.660	5.368	5.629	6.571	7.790	9.467	10.185	10.821	13.339	16.232	17.117	18.131	21.064	23.655	26.111	28.873	29.141	31.319	36.123	14
15	4.601	5.220	5.985	6.662	7.661	8.547	10.307	11.038	11.721	14.339	17.322	18.246	19.311	22.307	24.996	27.488	28.239	30.578	32.801	37.597	15
16	5.142	5.812	6.514	6.908	7.962	9.312	11.912	12.824	15.038	18.418	20.369	20.485	23.542	26.298	28.845	29.633	32.000	34.267	39.252	46.217	16
17	5.697	6.408	7.255	7.504	8.672	10.085	12.792	13.531	16.338	19.511	20.689	21.615	24.769	27.587	30.191	32.409	35.718	40.790	47.177	51.384	17
18	6.265	7.015	7.906	8.231	9.390	10.855	12.857	13.675	14.440	17.033	20.011	21.605	22.790	25.989	28.809	31.520	32.346	34.805	37.156	42.312	48.16
19	6.844	7.633	8.567	8.907	10.117	11.651	13.116	14.582	15.352	18.038	21.689	22.718	23.900	27.204	30.144	32.852	33.637	36.582	43.820	49.219	55.315
20	7.434	8.280	9.237	9.591	10.851	12.443	14.578	15.452	16.352	18.938	21.689	22.775	23.828	26.078	28.412	31.410	34.852	35.020	37.566	39.997	45.315
21	8.034	8.897	9.915	10.283	11.591	13.240	15.445	16.344	17.182	20.337	23.858	24.335	26.171	29.615	32.671	35.479	36.314	38.932	41.401	46.707	51
22	8.643	9.542	10.600	10.982	12.338	14.041	16.314	17.240	18.101	21.337	24.939	26.019	30.813	33.924	36.781	37.659	40.289	42.796	46.208	52.22	58
23	9.250	10.194	11.293	11.888	13.091	14.898	17.187	18.137	19.021	22.337	26.018	27.441	28.439	32.007	35.172	38.076	38.988	41.630	45.558	51.179	57.502
24	9.886	11.992	12.401	13.848	15.092	17.939	20.807	21.357	22.172	28.339	30.675	30.652	37.652	40.646	41.506	44.314	46.928	52.620	58.320	64.315	70
25	10.520	11.524	12.697	13.120	14.611	16.473	18.940	19.939	20.807	24.337	28.172	28.339	30.675	34.302	37.652	40.646	41.506	44.314	46.928	52.620	58
26	11.160	12.188	13.409	13.844	15.379	17.293	19.820	20.843	21.792	25.335	29.244	30.433	31.793	35.563	38.885	41.923	42.866	45.842	48.290	54.052	26
27	11.808	12.879	14.125	14.573	16.151	18.114	20.703	21.749	22.719	26.335	30.191	31.528	32.912	36.741	40.113	43.194	44.140	46.963	49.645	54.476	27
28	12.481	13.487	15.308	16.928	18.939	21.568	23.647	25.038	26.341	32.520	34.077	37.918	41.315	44.461	47.419	48.278	50.933	56.593	62.228	68.208	
29	13.121	14.256	15.574	16.047	17.700	19.768	22.475	23.567	24.577	28.339	32.481	33.711	35.139	38.087	42.557	45.722	46.693	49.888	52.358	58.302	64
30	13.787	14.933	16.306	16.791	18.493	20.599	23.364	24.478	25.508	29.335	33.530	36.800	36.230	40.256	43.773	46.979	47.962	50.932	53.672	59.703	60
40	20.706	22.184	23.638	24.433	25.509	29.031	32.345	33.660	34.872	39.335	44.195	45.616	47.389	51.805	55.759	59.342	60.436	63.191	66.766	73.402	40
40	20.706	22.184	23.638	24.433	25.509	29.031	32.345	33.660	34.872	39.335	44.195	45.616	47.389	51.805	55.759	59.342	60.436	63.191	66.766	73.402	40
50	27.901	29.707	31.664	32.357	34.188	41.449	42.942	44.311	45.723	50.334	55.327	56.981	58.164	62.187	67.505	72.613	76.154	79.490	86.561	90.501	95.521
60	35.535	37.495	39.699	40.482	43.188	46.459	50.641	52.294	53.809	58.305	65.227	66.981	68.972	74.397	79.082	83.298	84.580	88.379	91.932	99.607	100
70	43.255	45.492	47.893	48.758	51.739	55.329	59.898	61.346	64.334	65.069	71.577	79.716	85.527	90.511	95.523	96.388	100.125	104.215	112.311	122.517	70
80	51.171	53.539	56.213	57.153	60.301	64.278	67.145	71.251	73.334	78.120	86.130	91.577	96.588	101.680	106.099	112.321	116.321	124.839	137.208	90	
90	59.198	61.794	64.634	65.646	69.126	73.291	75.553	78.525	81.544	86.251	91.334	96.524	98.850	101.054	107.565	113.445	118.136	119.648	124.116	128.299	137.208
100	67.327	70.085	73.142	74.222	77.929	82.358	87.945	90.133	92.129	95.334	108.908	109.141	111.687	118.498	124.342	129.561	131.142	142.105	149.449	160.100	100

FINAL EXAMINATION

SEMESTER / SESSION : SEM I / 2014/2015
 COURSE : ENGINEERING
 MATHEMATICS V

PROGRAMME : 2 BEV / BEJ
 COURSE CODE : BEE 31702

Table 9


PERCENTAGE POINTS OF THE F DISTRIBUTION

The table gives the values of $F_{\alpha; \nu_1, \nu_2}$, the 100 α percentage point of the F distribution having ν_1 degrees of freedom in the numerator and ν_2 degrees of freedom in the denominator.

For each pair of values of ν_1 and ν_2 , $F_{\alpha; \nu_1, \nu_2}$ is tabulated for $\alpha = 0.05, 0.025, 0.01, 0.001$, the 0.025 values being bracketed.

The lower percentage points of the distribution may be obtained from the relation:-

$$F_{1-\alpha; \nu_1, \nu_2} = 1/F_{\alpha; \nu_2, \nu_1}$$



$$\text{e.g. } F_{.95; 12, 8} = 1/F_{.05; 8, 12} = 1/2.85 = 0.351$$

$\nu_1 \backslash \nu_2$	1	2	3	4	5	6	7	8	10	12	24	∞
1	161.4 (548)	199.5 (800)	215.7 (864)	224.6 (900)	230.2 (922)	234.0 (937)	236.8 (948)	238.9 (957)	241.9 (969)	243.9 (977)	249.0 (997)	254.3 (1018)
	4052	5000	5403	5625	5764	5859	5928	5981	6056	6106	6235	6366
	4053*	5000*	5404*	5625*	5764*	5859*	5929*	5981*	6056*	6107*	6235*	6366*
2	18.5 (38.5)	19.0 (39.0)	19.2 (39.2)	19.3 (39.3)	19.3 (39.3)	19.4 (39.4)	19.4 (39.4)	19.4 (39.4)	19.4 (39.4)	19.4 (39.4)	19.5 (39.5)	19.5 (39.5)
	98.5	99.0	99.2	99.3	99.3	99.4	99.4	99.4	99.4	99.4	99.5	99.5
	998.5	999.0	999.2	999.2	999.3	999.3	999.4	999.4	999.4	999.4	999.5	999.5
3	10.13 (17.4)	9.55 (16.0)	9.28 (15.4)	9.12 (15.1)	9.01 (14.9)	8.94 (14.7)	8.89 (14.6)	8.85 (14.5)	8.79 (14.4)	8.74 (14.3)	8.64 (14.1)	8.53 (13.9)
	34.1	30.8	29.5	28.7	28.2	27.9	27.7	27.5	27.2	27.1	26.6	26.1
	167.0	148.5	141.1	137.1	134.6	132.8	131.5	130.6	129.2	128.3	125.9	123.5
4	7.71 (12.22)	6.94 (10.65)	6.59 (9.98)	6.39 (9.60)	6.26 (9.36)	6.16 (9.20)	6.09 (9.07)	6.04 (8.98)	5.96 (8.84)	5.91 (8.75)	5.77 (8.51)	5.63 (8.26)
	21.2	18.0	16.7	16.0	15.5	15.2	15.0	14.8	14.5	14.4	13.9	13.5
	74.14	61.25	56.18	53.44	51.71	50.53	49.66	49.00	48.05	47.41	45.77	44.05
5	6.61 (10.01)	5.79 (8.43)	5.41 (7.76)	5.19 (7.39)	5.05 (7.15)	4.95 (6.98)	4.88 (6.85)	4.82 (6.76)	4.74 (6.62)	4.68 (6.52)	4.53 (6.28)	4.36 (6.02)
	16.26	13.27	12.06	11.39	10.97	10.67	10.46	10.29	10.05	9.89	9.47	9.02
	47.18	37.12	33.20	31.09	29.75	28.83	28.16	27.65	26.92	26.42	25.14	23.79
6	5.99 (8.81)	5.14 (7.26)	4.76 (6.60)	4.53 (6.23)	4.39 (5.99)	4.28 (5.82)	4.21 (5.70)	4.15 (5.60)	4.06 (5.46)	4.00 (5.37)	3.84 (5.12)	3.67 (4.85)
	13.74	10.92	9.78	9.15	8.75	8.47	8.26	8.10	7.87	7.72	7.31	6.88
	35.51	27.00	23.70	21.92	20.80	20.03	19.46	19.03	18.41	17.99	16.90	15.75
7	5.59 (8.07)	4.74 (6.54)	4.35 (5.89)	4.12 (5.52)	3.97 (5.29)	3.87 (5.12)	3.79 (4.99)	3.73 (4.90)	3.64 (4.76)	3.57 (4.67)	3.41 (4.42)	3.23 (4.14)
	12.25	9.55	8.45	7.85	7.46	7.19	6.99	6.84	6.62	6.47	6.07	5.65
	29.25	21.69	18.77	17.20	16.21	15.52	15.02	14.63	14.08	13.71	12.73	11.70
8	5.32 (7.57)	4.46 (6.06)	4.07 (5.42)	3.84 (5.05)	3.69 (4.82)	3.58 (4.65)	3.50 (4.53)	3.44 (4.43)	3.35 (4.30)	3.28 (4.20)	3.12 (3.95)	2.93 (3.67)
	11.26	8.65	7.59	7.01	6.63	6.37	6.18	6.03	5.81	5.67	5.28	4.86
	25.42	18.49	15.83	14.39	13.48	12.86	12.40	12.05	11.54	11.19	10.30	9.34
9	5.12 (7.21)	4.26 (5.71)	3.86 (5.08)	3.63 (4.72)	3.48 (4.48)	3.37 (4.32)	3.29 (4.20)	3.23 (4.10)	3.14 (3.96)	3.07 (3.87)	2.90 (3.61)	2.71 (3.33)
	10.56	8.02	6.99	6.42	6.06	5.80	5.61	5.47	5.26	5.11	4.73	4.31
	22.86	16.39	13.90	12.56	11.71	11.13	10.69	10.37	9.87	9.57	8.72	7.81
10	4.96 (6.94)	4.10 (5.46)	3.71 (4.83)	3.48 (4.47)	3.33 (4.24)	3.22 (4.07)	3.14 (3.95)	3.07 (3.85)	2.98 (3.72)	2.91 (3.62)	2.74 (3.37)	2.54 (3.08)
	10.04	7.56	6.55	5.99	5.64	5.39	5.20	5.06	4.85	4.71	4.33	3.91
	21.04	14.91	12.55	11.28	10.48	9.93	9.52	9.20	8.74	8.44	7.64	6.76
11	4.84 (6.72)	3.98 (5.26)	3.59 (4.63)	3.36 (4.28)	3.20 (4.04)	3.09 (3.88)	3.01 (3.76)	2.95 (3.66)	2.85 (3.53)	2.79 (3.43)	2.61 (3.17)	2.40 (2.88)
	9.65	7.21	6.22	5.67	5.32	5.07	4.89	4.74	4.54	4.40	4.02	3.60
	19.69	13.81	11.56	10.35	9.58	9.05	8.66	8.35	7.92	7.63	6.85	6.00
12	4.75 (6.55)	3.89 (5.10)	3.49 (4.47)	3.26 (4.12)	3.11 (3.89)	3.00 (3.73)	2.91 (3.61)	2.85 (3.51)	2.75 (3.37)	2.69 (3.28)	2.51 (3.02)	2.30 (2.72)
	9.33	6.93	5.95	5.41	5.06	4.82	4.64	4.50	4.30	4.16	3.78	3.36
	18.64	12.97	10.80	9.63	8.89	8.38	8.00	7.71	7.29	7.00	6.25	5.42