



**UNIVERSITI TUN HUSSEIN ONN MALAYSIA**

**FINAL EXAMINATION  
SEMESTER II  
SESSION 2016/2017**

COURSE NAME : ENGINEERING TECHNOLOGY  
MATHEMATICS II

COURSE CODE : BDU 11003

PROGRAMME CODE : BDC / BDM

EXAMINATION DATE : JUNE 2017

DURATION : 3 HOURS

INSTRUCTION : ANSWER ALL QUESTIONS IN  
**PART A AND THREE (3)  
QUESTIONS IN PART B.**

THIS QUESTION PAPER CONSISTS OF **SIX (6)** PAGES

**TERBUKA**

**CONFIDENTIAL**

**PART A**

**Q1** A periodic function  $f(x)$  is defined by

$$f(x) = \begin{cases} 1 & , -1 \leq x < 0, \\ 1-x & , 0 \leq x < 1, \end{cases}$$

and  $f(x) = f(x+2)$ .

(a) Sketch the graph of  $f(x)$  over  $-3 \leq x \leq 3$ .

(2 marks)

(b) Show that the Fourier coefficients corresponding to  $f(x)$  are

$$a_0 = \frac{3}{2}, \quad a_n = \begin{cases} \frac{2}{n^2\pi^2}, & n \text{ is odd} \\ 0 & , n \text{ is even} \end{cases} \quad \text{and} \quad b_n = \begin{cases} -\frac{1}{n\pi}, & n \text{ is odd} \\ \frac{1}{n\pi}, & n \text{ is even} \end{cases}.$$

(15 marks)

(c) Write the Fourier series of  $f(x)$  by giving your answers for the first three nonzero terms of  $a_n$  and  $b_n$ .

(3 marks)

**Q2** Given the heat equation

$$\frac{\partial u}{\partial t} = 0.4 \frac{\partial^2 u}{\partial x^2}, \quad \text{for } 0 < x < 1,$$

with the boundary conditions  $u(0, t) = e^{-t}$  and  $u(1, t) = t$  for  $t > 0$ . The initial condition is given by  $u(x, 0) = 2x$  for  $0 \leq x \leq 1$ .

(a) Draw a grid for this problem by taking  $\Delta x = h = 0.2$  and  $\Delta t = k = 0.25$ .

(3 marks)

(b) Write the given heat equation in finite-difference form where

$$\frac{\partial u_{i,j}}{\partial t} = c^2 \frac{\partial^2 u_{i,j}}{\partial x^2}$$

is approximated by

$$\frac{u_{i,j+1} - u_{i,j}}{k} = c^2 \frac{u_{i-1,j} - 2u_{i,j} + u_{i+1,j}}{h^2}.$$

(3 marks)

(c) From (b), find  $u(x, 0)$ ,  $u(x, 0.25)$  and  $u(x, 0.5)$ .

(14 marks)

## PART B

Q3 (a) Solve

$$(3x^2 - 2xy + e^y - ye^{-x}) dx + (2y - x^2 + e^{-x} + xe^y) dy = 0$$

with initial value  $y(0) = 1$ .

(11 marks)

(b) According to Newton's law of cooling, the rate at which a body cools is given by the equation

$$\frac{dT}{dt} = -k(T - T_s),$$

where  $T_s$  is the temperature of the surrounding medium,  $k$  is a constant and  $t$  is the time in minutes. If the body cools from  $100^\circ\text{C}$  to  $60^\circ\text{C}$  in 10 minutes with the surrounding temperature of  $20^\circ\text{C}$ , how long does it need for the body to cool from  $100^\circ\text{C}$  to  $25^\circ\text{C}$ .

(9 marks)

Q4 (a) By using an appropriate method, solve

$$y'' - 4y = 3x + e^{2x}$$

with  $y(0) = 0$  and  $y'(0) = 1$ .

(13 marks)

(b) A mass of 20.4 kg is suspended from a spring with a known spring constant of 29.4 N/m. The mass is set in motion from its equilibrium position with an upward velocity of 3.6m/s. The motion can be described in the differential equation

$$\ddot{x} + \frac{k}{m}x = 0$$

where  $m$  is the mass of the object and  $k$  is the spring constant.

(i) Determine the initial conditions.

(1 mark)

(ii) Find an equation for the position of the mass at any time  $t$ .

(6 marks)

TERBUKA

CONFIDENTIAL

**Q5** (a) Find the Laplace transform for each of the following function:

- (i)  $(2 + t^3)e^{-2t}$ .
- (ii)  $\sin(t - 2\pi)H(t - 2\pi)$ .
- (iii)  $\sin 3t \delta(2t - \pi)$ .

(10 marks)

(b) Consider the periodic function

$$f(t) = \begin{cases} t, & 0 \leq t < 1 \\ 1-t, & 1 \leq t < 2 \end{cases}$$

$$f(t) = f(t+2).$$

Sketch the graph of  $f(t)$  and find its Laplace transform.

$$\left[ \text{Hint: } \mathcal{L}[f(t)] = \frac{1}{1 - e^{-sT}} \int_0^T e^{-st} f(t) dt, \quad s > 0. \right]$$

(10 marks)

**Q6** (a) (i) Find the inverse Laplace transform of

$$\frac{s + 3}{s^2 - 6s + 13}$$

(ii) From (a)(i), find

$$\mathcal{L}^{-1} \left\{ \frac{(s + 3)e^{-\frac{1}{2}\pi s}}{s^2 - 6s + 13} \right\}$$

(8 marks)

(b) (i) Express

$$\frac{1}{(s - 1)(s - 2)^2}$$

in partial fractions and show that

$$\mathcal{L}^{-1} \left\{ \frac{1}{(s - 1)(s - 2)^2} \right\} = e^t - e^{2t} + te^{2t}.$$

(ii) Use the result in (i) to solve the differential equation

$$y' - y = te^{2t}$$

which satisfies the initial condition of  $y(0) = 1$ .

(12 marks)

**-END OF QUESTION-**

**FINAL EXAMINATION**

SEMESTER/SESSION: SEM II/2016/2017  
 COURSE NAME : ENGINEERING TECHNOLOGY  
 MATHEMATICS II

PROGRAMME : BDC/BDM  
 COURSE CODE: BDU 11003

**Formulae**

**Characteristic Equation and General Solution**

Case	Roots of the Characteristic Equation	General Solution
1	$m_1$ and $m_2$ ; real and distinct	$y = Ae^{m_1x} + Be^{m_2x}$
2	$m_1 = m_2 = m$ ; real and equal	$y = (A + Bx)e^{mx}$
3	$m = \alpha \pm i\beta$ ; imaginary	$y = e^{\alpha x} (A \cos \beta x + B \sin \beta x)$

**Particular Integral of  $ay'' + by' + cy = f(x)$  : Method of Undetermined Coefficients**

$f(x)$	$y_p(x)$
$P_n(x) = A_n x^n + \dots + A_1 x + A_0$	$x^r (B_n x^n + \dots + B_1 x + B_0)$
$Ce^{\alpha x}$	$x^r (Pe^{\alpha x})$
$C \cos \beta x$ or $C \sin \beta x$	$x^r (p \cos \beta x + q \sin \beta x)$

**Particular Integral of  $ay'' + by' + cy = f(x)$  : Method of Variation of Parameters**

Wronskian	Parameter	Solution
$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$	$u_1 = -\int \frac{y_2 f(x)}{W} dx, \quad u_2 = \int \frac{y_1 f(x)}{W} dx$	$y_p = u_1 y_1 + u_2 y_2$

**TERBUKA**

**CONFIDENTIAL**



**FINAL EXAMINATION**

SEMESTER/SESSION: SEM II/2016/2017  
 COURSE NAME : ENGINEERING TECHNOLOGY  
 MATHEMATICS II

PROGRAMME : BDC/BDM  
 COURSE CODE: BDU 11003

**Laplace Transforms**

$$\mathcal{L}\{f(t)\} = \int_0^{\infty} f(t)e^{-st} dt = F(s)$$

$f(t)$	$F(s)$	$f(t)$	$F(s)$
$a$	$\frac{a}{s}$	$H(t-a)$	$\frac{e^{-as}}{s}$
$t^n, n=1, 2, 3, \dots$	$\frac{n!}{s^{n+1}}$	$f(t-a)H(t-a)$	$e^{-as}F(s)$
$e^{at}$	$\frac{1}{s-a}$	$\delta(t-a)$	$e^{-as}$
$\sin at$	$\frac{a}{s^2+a^2}$	$f(t)\delta(t-a)$	$e^{-as}f(a)$
$\cos at$	$\frac{s}{s^2+a^2}$	$\int_0^t f(u)g(t-u)du$	$F(s).G(s)$
$\sinh at$	$\frac{a}{s^2-a^2}$	$y(t)$	$Y(s)$
$\cosh at$	$\frac{s}{s^2-a^2}$	$\dot{y}(t)$	$sY(s) - y(0)$
$e^{at}f(t)$	$F(s-a)$	$\ddot{y}(t)$	$s^2Y(s) - sy(0) - \dot{y}(0)$
$t^n f(t), n=1, 2, 3, \dots$	$(-1)^n \frac{d^n F(s)}{ds^n}$		

**Periodic Function for Laplace transform :**  $\mathcal{L}\{f(t)\} = \frac{1}{1-e^{-sT}} \int_0^T e^{-st} f(t) dt, \quad s > 0.$

**Fourier Series**

$$f(x) = \frac{a_0}{2} + \sum_{n=1}^{\infty} \left[ a_n \cos \frac{n\pi x}{L} + b_n \sin \frac{n\pi x}{L} \right]$$

where

$$a_0 = \frac{1}{L} \int_{-L}^L f(x) dx$$

$$a_n = \frac{1}{L} \int_{-L}^L f(x) \cos \frac{n\pi x}{L} dx$$

$$b_n = \frac{1}{L} \int_{-L}^L f(x) \sin \frac{n\pi x}{L} dx$$

**TERBUKA**